

AD-A147 061

STICKING PROBABILITY OF H2 AND HD ON NOBLE METAL
SURFACES(U) CORNELL UNIV ITHACA NY LAB OF ATOMIC AND
SOLID STATE PHYSICS M D STILES ET AL. 1980 TR-6

1/1


UNCLASSIFIED

N00014-80-C-0489

F/G 7/4

NL





12

OFFICE OF NAVAL RESEARCH

Research Contract No. N00014-80-C-0489

TECHNICAL REPORT No. 6

STICKING PROBABILITY OF H_2 AND HD ON NOBLE METAL SURFACES

by

M.D. Stiles and J.W. Wilkins

Prepared for Publication

in

Physical Review Letters

Laboratory of Atomic and Solid State Physics
Cornell University
Ithaca, NY 14853

Reproduction in whole or part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited.

AD-A147 061

DTIC FILE COPY

DTIC
ELECTE
NOV 2 1984
S D D

84 10 26 017

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability	
Dist	Availability, of Special
A/1	



Sticking Probability of H_2 and HD on Noble Metal Surfaces

M. D. Stiles and J. W. Wilkins

Laboratory of Atomic and Solid State Physics
Cornell University
Ithaca, NY 14853

One-phonon distorted-wave Born approximation calculations of the energy-dependent phonon-assisted sticking coefficients of H_2 and HD at smooth metal surfaces yield peaks in the sticking coefficient as a function of energy due to selective adsorption resonances, the peaks being strong for HD and weak for H_2 . Furthermore the calculation does not explain the observed¹ dependence of the sticking coefficient of H_2 on the rotational populations.

PACS numbers: 82.65.My., 68.30.+z.

END

FILMED

11-84

DTIC